


Types of Forces

- Force balance





$$\sum_i \vec{F}_i = m\vec{a}$$

Energy and laws of conservation




- Work
 - Energy
 - Power
 - Rotational Motion
 - Many-Particle Interactions
-

Work



$$\Delta W = \vec{F} \Delta \vec{r} \quad W = \int_{r_1}^{r_2} \vec{F} d\vec{r} \quad \left[\frac{\text{kg m}^2}{\text{s}^2}, \text{Nm}, J \right]$$



Energy



■ Kinetic Energy






$$E_k = \frac{mv^2}{2} = \frac{p^2}{2m}$$

Kinetic energy is the energy of an object due to its motion

Energy



■ Potential Energy



$$E_p = mgh$$


$$E_p = \frac{k\Delta x^2}{2}$$

Potential energy is the energy of an object due to its location

Energy



Thermal energy



Chemical energy

In the process of photosynthesis, plants convert radiant energy from the sun into chemical energy in the form of glucose - or sugar.



Nuclear energy



Electromagnetic energy

Power




$$N = \frac{\Delta W}{\Delta t} \quad \left[\frac{J}{s}, \text{ watt} \right]$$

Power is defined as the amount of work performed per unit time.



Rotational Motion

■ Definitions




Angular position: $\vec{\theta}(t)$

Angular velocity: $\vec{\omega}(t) = \dot{\vec{\theta}}(t)$ $[s^{-1}]$

Frequency: $f = \frac{\omega}{2\pi}$ $[s^{-1}]$

Angular acceleration: $\vec{\epsilon}(t) = \dot{\vec{\omega}}(t) = \ddot{\vec{\theta}}(t)$ $[s^{-2}]$






Tangential velocity: $\vec{v}(t) = [\vec{\omega}(t) \times \vec{r}(t)]$ $\left[\frac{m}{s} \right]$

Tangential acceleration: $\vec{a}(t) = \frac{d}{dt} [\vec{\omega}(t) \times \vec{r}(t)] =$
 $= [\vec{\epsilon}(t) \times \vec{r}(t)] + [\vec{\omega}(t) \times [\vec{\omega}(t) \times \vec{r}(t)]]$ $\left[\frac{m}{s^2} \right]$




Rotational Motion

■ Angular momentum





Angular momentum: $\vec{L} = [\vec{r} \times \vec{p}]$






Rotational Motion

■ Torque



Torque: $\vec{M} = [\vec{r} \times \vec{F}]$ $[Nm]$



Rotational Motion

□ Moment of inertia I [kgm^2]


$$I = \sum_i m_i r_i^2 \quad [\text{kg m}^2]$$

$$I = \int_V r^2 dm \quad [\text{kg m}^2]$$


$$\vec{L} = J \vec{\omega}$$

$$\vec{M} = J \vec{\epsilon}$$


$$T = \frac{I\omega^2}{2}$$




Bowling ball (solid)



Cylinder



Cylinder





Beachball (shell)



Cylinder

Inertia of some selected shapes

Rotational Motion





$$\vec{L} = J \vec{\omega}$$

$$\vec{M} = J \vec{\epsilon}$$

$$T = \frac{I\omega^2}{2}$$

Rotational Motion





$$\vec{L} = J \vec{\omega}$$

$$\vec{M} = J \vec{\epsilon}$$

$$T = \frac{I\omega^2}{2}$$

Rotational Motion





$$\vec{L} = J \vec{\omega}$$

$$\vec{M} = J \vec{\varepsilon}$$

$$T = \frac{I\omega^2}{2}$$

Rotational Motion




$$\vec{L} = J \vec{\omega}$$

$$\vec{M} = J \vec{\epsilon}$$

$$T = \frac{I \omega^2}{2}$$

Rotational Motion




$$\vec{L} = \hat{J} \vec{\omega}$$

$$\vec{M} = \hat{J} \vec{\epsilon}$$

- Inertia tensor

$$J = \begin{bmatrix} I_{xx} & -I_{xy} & I_{xz} \\ I_{xy} & I_{yy} & -I_{yz} \\ -I_{xz} & I_{yz} & I_{zz} \end{bmatrix}$$




Kinetic energy:

$$T = \frac{1}{2} \vec{\omega} \hat{J} \vec{\omega}$$



Angular Momentum of a Rigid Body



Axis of rotation



$$\vec{L} = \sum_i \vec{l}_i = \sum_i [\vec{r}_i \times m_i \vec{v}] = \sum_i m_i [\vec{r}_i \times [\vec{\omega} \times \vec{r}_i]]$$

$$= \sum_i m_i \begin{bmatrix} 0 & -z_i & y_i \\ z_i & 0 & -x_i \\ -y_i & x_i & 0 \end{bmatrix} \begin{bmatrix} 0 & -\omega_z & \omega_y \\ \omega_z & 0 & -\omega_x \\ -\omega_y & \omega_x & 0 \end{bmatrix} \begin{bmatrix} x_i \\ y_i \\ z_i \end{bmatrix}$$

$$= \sum_i m_i \begin{bmatrix} 0 & -z_i & y_i \\ z_i & 0 & -x_i \\ -y_i & x_i & 0 \end{bmatrix} \begin{bmatrix} z_i \omega_y - y_i \omega_z \\ x_i \omega_z - z_i \omega_x \\ y_i \omega_x - x_i \omega_y \end{bmatrix} = \sum_i m_i \begin{bmatrix} \omega_x (y_i^2 + z_i^2) - \omega_y x_i y_i - \omega_z x_i z_i \\ -\omega_x x_i y_i + \omega_y (x_i^2 + z_i^2) - \omega_z y_i z_i \\ -\omega_x z_i x_i - \omega_y y_i z_i + \omega_z (x_i^2 + y_i^2) \end{bmatrix}$$

$$= \begin{bmatrix} \left(\sum_i m_i (y_i^2 + z_i^2) \right) \omega_x + \left(-\sum_i m_i x_i y_i \right) \omega_y + \left(-\sum_i m_i x_i z_i \right) \omega_z \\ \left(-\sum_i m_i x_i y_i \right) \omega_x + \left(\sum_i m_i (x_i^2 + z_i^2) \right) \omega_y + \left(-\sum_i m_i y_i z_i \right) \omega_z \\ \left(-\sum_i m_i x_i z_i \right) \omega_x + \left(-\sum_i m_i z_i y_i \right) \omega_y + \left(\sum_i m_i (x_i^2 + y_i^2) \right) \omega_z \end{bmatrix} = \begin{bmatrix} \left(\sum_i m_i (y_i^2 + z_i^2) \right) & \left(-\sum_i m_i x_i y_i \right) & \left(-\sum_i m_i x_i z_i \right) \\ \left(-\sum_i m_i x_i y_i \right) & \left(\sum_i m_i (x_i^2 + z_i^2) \right) & \left(-\sum_i m_i y_i z_i \right) \\ \left(-\sum_i m_i x_i z_i \right) & \left(-\sum_i m_i z_i y_i \right) & \left(\sum_i m_i (x_i^2 + y_i^2) \right) \end{bmatrix} \begin{bmatrix} \omega_x \\ \omega_y \\ \omega_z \end{bmatrix}$$

Physical pendulum



$$\vec{F} = -k\Delta \vec{x}$$

$$\vec{M} = -k\Delta \vec{\theta}$$

$$\vec{M} = \hat{J} \ddot{\vec{\theta}}$$

$$J \ddot{\theta} + k\theta = 0$$

$$\ddot{\theta} + \omega^2 \theta = 0$$

Physical pendulum




Rigid Body Roller Coaster

myPhysicsLab.com

English [previous](#) [next](#)

Sim Graph Time Graph Multi Graph



path

gravity

damping

elasticity

show forces

show energy

show clock

pan-zoom

time step


time rate

Diff Eq Solver

potential energy offset


background

terminal



Many-Particle Interactions

- For an N particles system, in equilibrium



$$\vec{P} = \sum_i \vec{p}_i = \sum_i m_i \vec{v}_i$$

$$\vec{L} = \sum_i \vec{r}_i \times \vec{p}_i = \sum_i \vec{r}_i \times m \vec{v}_i$$

$$\sum_i \vec{F}_i = 0$$

resultant force = 0

$$\sum_i \vec{r}_i \times \vec{F}_i = 0$$

resultant moment (torque) = 0

Conservation laws




$$\sum_i m_i = \text{const}$$

$$\sum_i E_i = \text{const}$$

$$\sum_i \vec{P}_i = \text{const}$$


$$\sum_i \vec{L}_i = \text{const}$$

Projectiles





- Topics
 - The gravity-only model
 - Aerodynamic drag
 - Laminar and turbulent flow
 - Wind effects
 - Spin effects
 - Details on specific types of projectiles including bullets, cannonballs, and arrows
-

Projectiles



- The gravity-only model



Equation of motion:

$$\frac{d\vec{v}}{dt} = \vec{g} \quad \text{or} \quad \frac{d^2\vec{r}}{dt^2} = \vec{g}$$

Solution:

$$\vec{v} = \vec{v}_0 + \vec{g}t$$

$$\vec{r} = \vec{r}_0 + \vec{v}_0t + \frac{1}{2}\vec{g}t^2$$

or


$$v_y = v_{y0} - gt,$$

$$v_x = v_{x0}$$

$$y = y_0 + v_{y0}t - \frac{1}{2}gt^2,$$

$$x = x_0 + v_{x0}t,$$

Projectiles



- The gravity-only model



Equation of motion:

$$\frac{d\vec{v}}{dt} = \vec{g} \quad \text{or} \quad \frac{d^2\vec{r}}{dt^2} = \vec{g}$$

Solution:

$$\vec{v} = \vec{v}_0 + \vec{g}t$$

$$\vec{r} = \vec{r}_0 + \vec{v}_0t + \frac{1}{2}\vec{g}t^2$$

or


$$v_y = v_{y0} - gt,$$

$$v_x = v_{x0}$$


$$y = y_0 + v_{y0}t - \frac{1}{2}gt^2,$$

$$x = x_0 + v_{x0}t,$$

Projectiles



■ The gravity-only model



Equation of motion:

$$\frac{d\vec{v}}{dt} = \vec{g} \quad \text{or} \quad \frac{d^2\vec{r}}{dt^2} = \vec{g}$$

Solution:

$$\vec{v} = \vec{v}_0 + \vec{g}t$$

$$\vec{r} = \vec{r}_0 + \vec{v}_0t + \frac{1}{2}\vec{g}t^2$$

or

$$v_y = v_{y0} - gt,$$

$$v_x = v_{x0}$$


$$y = y_0 + v_{y0}t - \frac{1}{2}gt^2,$$

$$x = x_0 + v_{x0}t,$$




Projectiles

- The gravity-only model
 - Golf Game



Projectiles



- The gravity-only model
 - Summary
 - The only force on the projectile is due to gravity, which acts in the vertical.
 - The motion in the three coordinate directions is independent.
 - The projectile trajectory is independent of mass and projectile geometry.
 - The velocity in the x- and y-directions is constant over the entire trajectory and is equal to the initial velocities in the x- and y-direction.
 - The shape of the projectile trajectory is a parabola.
-

Projectiles




- **Aerodynamic drag**
 - **Basic Concepts**



$$\vec{F}_{\text{Drag}} = \vec{F}_{D, \text{pressure}} + \vec{F}_{D, \text{Friction}}$$

Total drag Pressure drag Friction drag (or skin drag)

Projectiles




- **Aerodynamic drag**
 - **Drag force**

$$F_d = C_d A \rho \frac{V^2}{2}$$

Drag force → F_d
 Drag coefficient → C_d
 Effective area → A
 Density of the fluid → ρ
 Velocity → V

Shape	Picture	C_D
Square flat plate		1.17
Cube		1.05–1.07
Rotated cube		0.8–0.81
Solid hemisphere		0.42
60-degree cone		0.5
Sphere		0.4–0.47
2:1 Ellipsoid		0.27
Hollow hemisphere		1.4
Hollow hemisphere		0.38–0.4


Projectiles



- **Aerodynamic drag**
 - **Drag Coefficient**
 - **Laminar and Turbulent Flow**


$$C_d = C_d(\text{Re})$$

$$\text{Re} = \frac{\rho v L}{\mu}$$



The drag coefficient of a sphere as a function of Reynolds number

Projectiles



- **Aerodynamic drag**
 - **Drag Coefficient**
 - **Laminar and Turbulent Flow**


$$C_D = C_D(\text{Re})$$

$$\text{Re} = \frac{\rho v L}{\mu}$$

Laminar and Turbulent Drag Coefficients

Shape	Laminar C_D	Turbulent C_D
Sphere	0.4–0.47	0.2
2:1 Ellipsoid	0.27	0.13
Circular cylinder	1.2	0.3
2:1 Elliptical cylinder	0.6	0.2

Projectiles




- **Aerodynamic drag**
 - **Drag Coefficient**
 - **Altitude Effects on Density**

Values of Air Density As a Function of Altitude


Altitude (m)	Altitude (ft)	Density (kg/m ³)	Density (slug/ft ³)
0.0	0.0	1.225	0.00238
305	1000	1.189	0.00231
610	2000	1.154	0.00224
914	3000	1.121	0.00218
1219	4000	1.088	0.00211
1524	5000	1.055	0.00205
2134	7000	0.992	0.00192
3048	10,000	0.905	0.00176

Projectiles



■ Aerodynamic drag

● Equations of motion



$$\vec{F} = m \vec{g} - |\vec{F}_D| \frac{\vec{v}}{|\vec{v}|}$$

$$F_d = C_d A \rho \frac{V^2}{2}$$



$$\vec{r} = \vec{g} - \frac{|\vec{F}_D|}{m} \frac{\vec{r}}{|\dot{\vec{r}}|}$$




$$\frac{dv_x}{dt} = a_x = -\frac{F_D v_x}{mv}$$


$$\frac{dv_y}{dt} = a_y = -\frac{F_D v_y}{mv}$$

$$\frac{dv_z}{dt} = a_z = -g - \frac{F_D v_z}{mv}$$


Projectiles




- **Aerodynamic drag**
 - **Three beachballs 2**




Projectiles




- **Aerodynamic drag**
 - **Golf Game Version 2**



Projectiles



- **Aerodynamic drag**
 - **Bowling balls**



Projectiles



■ Aerodynamic drag

● Summary


- Drag force acts in the opposite direction to the velocity. The magnitude of the drag force is proportional to the square of the velocity.
 - The three components of motion are coupled when drag is taken into account.
 - The drag force is a function of the projectile geometry.
 - The acceleration due to drag is inversely proportional to the mass of the projectile.
 - The drag on an object is proportional to the density of the fluid in which it is traveling.
-

Projectiles



■ Wind Effect

- Equations of motion



Apparent velocity is the vector sum of the projectile velocity and wind velocity.

$$\vec{F} = m \vec{g} - |\vec{F}_D| \frac{\vec{v}}{|\vec{v}|}$$



$$\vec{r} = \vec{g} - \frac{|\vec{F}_D|}{m} \frac{\vec{r}}{|\vec{r}|}$$



$$\frac{dv_x}{dt} = a_x = -\frac{F_D v_x}{mv}$$

$$\frac{dv_y}{dt} = a_y = -\frac{F_D v_y}{mv}$$


$$\frac{dv_z}{dt} = a_z = -g - \frac{F_D v_z}{mv}$$

$$F_d = C_d A \rho \frac{V^2}{2}$$

Projectiles



- **Wind Effect**
 - **Golf Game Version 3**




The effects of headwind or tailwind on a golf ball trajectory

...Java_Code\Chapter05_Projectile\GolfGame3.java (<https://ae.onliu.se/tsbk03>)

Projectiles



■ Wind Effect



Projectiles



■ Wind Effect



● Summary

- The presence of wind changes the apparent velocity seen by the projectile in flight. A headwind will increase the apparent velocity. A tailwind will decrease it.



 - The wind velocity affects the drag force in all three coordinate directions even if the wind velocities themselves are only in the x- and z-planes.
-

Projectiles

- Projectiles
 - Spin Effect



Bernoulli's equation




Bernoulli's equation: $p + \frac{1}{2} \rho v^2 + \rho g h = \text{const}$


pressure → p
 the fluid density → ρ
 velocity → v
 altitude → h

$\Rightarrow p + \frac{1}{2} \rho v^2 = \text{const}$
 ($h = \text{const}$)

Projectiles



- Projectiles
 - Spin Effect



⇒ **A spinning object generates lift**

Bernoulli's equation: $p + \frac{1}{2} \rho v^2 + \rho gh = const$

pressure → p , the fluid density → ρ , velocity → v , altitude → h

⇒ $p + \frac{1}{2} \rho v^2 = const$
($h = const$)


Projectiles



Spin Effect

- Magnus force

$$\vec{F}_M = C_L \rho \frac{v^2}{2} A \frac{[\vec{\omega} \times \vec{v}]}{||\vec{\omega} \times \vec{v}||}$$



The Magnus force lift coefficient

For a sphere: $C_L = \frac{r\omega}{v}$

For a cylinder: $C_L = \frac{2\pi r\omega}{v}$

- Equation of motion

$$\vec{F} = m \vec{g} - |\vec{F}_D| \frac{\vec{v}}{|\vec{v}|} + C_L \rho \frac{v^2}{2} \frac{[\vec{\omega} \times \vec{v}]}{||\vec{\omega} \times \vec{v}||}$$




$$\vec{\ddot{r}} = \vec{g} - \frac{|\vec{F}_D|}{m} \frac{\vec{r}}{|\dot{\vec{r}}|} + C_L \rho \frac{|\dot{\vec{r}}|^2}{2m} \frac{[\dot{\vec{\theta}} \times \dot{\vec{r}}]}{||\dot{\vec{\theta}} \times \dot{\vec{r}}||}$$

Projectiles




- Spin Effect
 - Magnus force demo



Projectiles



- Spin Effect
 - Golf Game Version 4



The effect of spin on golf ball flight

...Java_Code\Chapter05_Projectile\GolfGame4.java (<https://ae.onliu.se/tsbk03>)